

## 5. Random Vectors

- probabilities
- characteristic function
- cross correlation, cross covariance
- Gaussian random vectors
- functions of random vectors

# Random vectors

we denote  $\mathbf{X}$  a random vector

$\mathbf{X}$  is a function that maps each outcome  $\zeta$  to a vector of real numbers

an  $n$ -dimensional random variable has  $n$  components:

$$\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}$$

also called a *multivariate* or *multiple* random variable

# Probabilities

## Joint CDF

$$F(\mathbf{x}) \triangleq F_{\mathbf{X}}(x_1, x_2, \dots, x_n) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n)$$

## Joint PMF

$$p(\mathbf{x}) \triangleq p_{\mathbf{X}}(x_1, x_2, \dots, x_n) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

## Joint PDF

$$f(\mathbf{x}) \triangleq f_{\mathbf{X}}(x_1, x_2, \dots, x_n) = \frac{\partial^n}{\partial x_1 \dots \partial x_n} F(\mathbf{x})$$

## Marginal PMF

$$p_{X_j}(x_j) = P(X_j = x_j) = \sum_{x_1} \cdots \sum_{x_{j-1}} \sum_{x_{j+1}} \cdots \sum_{x_n} p_{\mathbf{X}}(x_1, x_2, \dots, x_n)$$

## Marginal PDF

$$f_{X_j}(x_j) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{\mathbf{X}}(x_1, x_2, \dots, x_n) dx_1 \cdots dx_{j-1} dx_{j+1} \cdots dx_n$$

**Conditional PDF:** the PDF of  $X_n$  given  $X_1, \dots, X_{n-1}$  is

$$f(x_n | x_1, \dots, x_{n-1}) = \frac{f_{\mathbf{X}}(x_1, \dots, x_n)}{f_{X_1, \dots, X_{n-1}}(x_1, \dots, x_{n-1})}$$

# Characteristic Function

the characteristic function of an  $n$ -dimensional RV is defined by

$$\begin{aligned}\Phi(\boldsymbol{\omega}) = \Phi(\omega_1, \dots, \omega_n) &= \mathbf{E}[e^{j(\omega_1 X_1 + \dots + \omega_n X_n)}] \\ &= \int_{\mathbf{x}} e^{j\boldsymbol{\omega}^T \mathbf{x}} f(\mathbf{x}) d\mathbf{x}\end{aligned}$$

where

$$\boldsymbol{\omega} = \begin{bmatrix} \omega_1 \\ \omega_2 \\ \vdots \\ \omega_n \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$\Phi(\boldsymbol{\omega})$  is the  $n$ -dimensional Fourier transform of  $f(\mathbf{x})$

# Independence

the random variables  $X_1, \dots, X_n$  are **independent** if

the joint pdf (or pmf) is equal to the product of their marginal's

## Discrete

$$p_{\mathbf{X}}(x_1, \dots, x_n) = p_{X_1}(x_1) \cdots p_{X_n}(x_n)$$

## Continuous

$$f_{\mathbf{X}}(x_1, \dots, x_n) = f_{X_1}(x_1) \cdots f_{X_n}(x_n)$$

we can specify an RV by the characteristic function in place of the pdf,

$X_1, \dots, X_n$  are *independent* if

$$\Phi(\omega) = \Phi_1(\omega_1) \cdots \Phi_n(\omega_n)$$

## Example: White noise signal in communication

the  $n$  samples  $X_1, \dots, X_n$  of a noise signal have the joint pdf:

$$f_{\mathbf{X}}(x_1, \dots, x_n) = \frac{e^{-(x_1^2 + \dots + x_n^2)/2}}{(2\pi)^{n/2}} \quad \text{for all } x_1, \dots, x_n$$

the joint pdf is the  $n$ -product of one-dimensional Gaussian pdf's

thus,  $X_1, \dots, X_n$  are independent Gaussian random variables

# Expected Values

the expected value of a function

$$g(\mathbf{X}) = g(X_1, \dots, X_n)$$

of a vector random variable  $\mathbf{X}$  is defined by

$$\mathbf{E}[g(\mathbf{X})] = \int_{\mathbf{x}} g(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} \quad \text{Continuous}$$

$$\mathbf{E}[g(\mathbf{X})] = \sum_{\mathbf{x}} g(\mathbf{x}) p(\mathbf{x}) \quad \text{Discrete}$$

**Mean vector**

$$\boldsymbol{\mu} = \mathbf{E}[\mathbf{X}] = \mathbf{E} \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \triangleq \begin{bmatrix} \mathbf{E}[X_1] \\ \mathbf{E}[X_2] \\ \vdots \\ \mathbf{E}[X_n] \end{bmatrix}$$



# Correlation and Covariance matrices

**Correlation matrix** has the second moments of  $\mathbf{X}$  as its entries:

$$\mathbf{R} \triangleq \mathbf{E}[\mathbf{X}\mathbf{X}^T] = \begin{bmatrix} \mathbf{E}[X_1X_1] & \mathbf{E}[X_1X_2] & \cdots & \mathbf{E}[X_1X_n] \\ \mathbf{E}[X_2X_1] & \mathbf{E}[X_2X_2] & \cdots & \mathbf{E}[X_2X_n] \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{E}[X_nX_1] & \mathbf{E}[X_nX_2] & \cdots & \mathbf{E}[X_nX_n] \end{bmatrix}$$

with

$$\mathbf{R}_{ij} = \mathbf{E}[X_iX_j]$$

**Covariance matrix** has the second-order central moments as its entries:

$$\mathbf{C} \triangleq \mathbf{E}[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T]$$

with

$$\mathbf{C}_{ij} = \mathbf{cov}(X_i, X_j) = \mathbf{E}[(X_i - \mu_i)(X_j - \mu_j)]$$

# Symmetric matrix

$A \in \mathbf{R}^{n \times n}$  is called *symmetric* if  $A = A^T$

**Facts:** if  $A$  is symmetric

- all eigenvalues of  $A$  are real
- all eigenvectors of  $A$  are orthogonal
- $A$  admits a decomposition

$$A = UDU^T$$

where  $U^T U = U U^T = I$  ( $U$  is unitary) and  $D$  is diagonal

(of course, the diagonals of  $D$  are eigenvalues of  $A$ )

# Unitary matrix

a matrix  $U \in \mathbf{R}^{n \times n}$  is called **unitary** if

$$U^T U = U U^T = I$$

**example:**  $\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

## Facts:

- a real unitary matrix is also called **orthogonal**
- a unitary matrix is always invertible and  $U^{-1} = U^T$
- columns vectors of  $U$  are mutually orthogonal
- norm is preserved under a unitary transformation:

$$y = Ux \implies \|y\| = \|x\|$$

# Positive definite matrix

a symmetric matrix  $A$  is **positive semidefinite**, written as  $A \succeq 0$  if

$$x^T A x \geq 0, \quad \forall x \in \mathbf{R}^n$$

and **positive definite**, written as  $A \succ 0$  if

$$x^T A x > 0, \quad \text{for all } \textit{nonzero } x \in \mathbf{R}^n$$

**Facts:**  $A \succeq 0$  if and only if

- all eigenvalues of  $A$  are non-negative
- all principle minors of  $A$  are non-negative

**example:**  $A = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} \succeq 0$  because

$$\begin{aligned} x^T A x &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\ &= x_1^2 + 2x_2^2 - 2x_1x_2 \\ &= (x_1 - x_2)^2 + x_2^2 \geq 0 \end{aligned}$$

or we can check from

- eigenvalues of  $A$  are 0.38 and 2.61 (real and positive)
- the principle minors are 1 and  $\begin{vmatrix} 1 & -1 \\ -1 & 2 \end{vmatrix} = 1$  (all positive)

note:  $A \succeq 0$  does not mean all entries of  $A$  are positive!

# Properties of correlation and covariance matrices

let  $\mathbf{X}$  be a (real)  $n$ -dimensional random vector with mean  $\mu$

## Facts:

- $\mathbf{R}$  and  $\mathbf{C}$  are  $n \times n$  symmetric matrices
- $\mathbf{R}$  and  $\mathbf{C}$  are positive semidefinite
- If  $X_1, \dots, X_n$  are independent, then  $\mathbf{C}$  is diagonal
- the diagonals of  $\mathbf{C}$  are given by the variances of  $X_k$
- if  $\mathbf{X}$  has zero mean, then  $\mathbf{R} = \mathbf{C}$
- $\mathbf{C} = \mathbf{R} - \mu\mu^T$

# Cross Correlation and Cross Covariance

let  $\mathbf{X}, \mathbf{Y}$  be vector random variables with means  $\mu_X, \mu_Y$  respectively

## Cross Correlation

$$\mathbf{cor}(\mathbf{X}, \mathbf{Y}) = \mathbf{E}[\mathbf{X}\mathbf{Y}^T]$$

if  $\mathbf{cor}(\mathbf{X}, \mathbf{Y}) = 0$  then  $\mathbf{X}$  and  $\mathbf{Y}$  are said to be **orthogonal**

## Cross Covariance

$$\begin{aligned}\mathbf{cov}(\mathbf{X}, \mathbf{Y}) &= \mathbf{E}[(\mathbf{X} - \mu_X)(\mathbf{Y} - \mu_Y)^T] \\ &= \mathbf{cor}(\mathbf{X}, \mathbf{Y}) - \mu_X\mu_Y^T\end{aligned}$$

if  $\mathbf{cov}(\mathbf{X}, \mathbf{Y}) = 0$  then  $\mathbf{X}$  and  $\mathbf{Y}$  are said to be **uncorrelated**

# Affine transformation

let  $\mathbf{Y}$  be an affine transformation of  $\mathbf{X}$ :

$$\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$$

where  $\mathbf{A}$  and  $\mathbf{b}$  are deterministic matrices

- $\mu_Y = \mathbf{A}\mu_X + \mathbf{b}$

$$\mu_Y = \mathbf{E}[\mathbf{A}\mathbf{X} + \mathbf{b}] = \mathbf{A} \mathbf{E}[\mathbf{X}] + \mathbf{E}[\mathbf{b}] = \mathbf{A}\mu_X + \mathbf{b}$$


- $\mathbf{C}_Y = \mathbf{A}\mathbf{C}_X\mathbf{A}^T$

$$\begin{aligned}\mathbf{C}_Y &= \mathbf{E}[(\mathbf{Y} - \mu_Y)(\mathbf{Y} - \mu_Y)^T] = \mathbf{E}[(\mathbf{A}(\mathbf{X} - \mu_X))(\mathbf{A}(\mathbf{X} - \mu_X))^T] \\ &= \mathbf{A} \mathbf{E}[(\mathbf{X} - \mu_X)(\mathbf{X} - \mu_X)^T] \mathbf{A}^T = \mathbf{A}\mathbf{C}_X\mathbf{A}^T\end{aligned}$$



# Diagonalization of covariance matrix

suppose a random vector  $\mathbf{Y}$  is obtained via a linear transformation of  $\mathbf{X}$



- the covariance matrices of  $\mathbf{X}$ ,  $\mathbf{Y}$  are  $\mathbf{C}_X$ ,  $\mathbf{C}_Y$  respectively
- $\mathbf{A}$  may represent linear filter, system gain, etc.
- the covariance of  $\mathbf{Y}$  is  $\mathbf{C}_Y = \mathbf{A}\mathbf{C}_X\mathbf{A}^T$

**Problem:** choose  $\mathbf{A}$  such that  $\mathbf{C}_Y$  becomes 'diagonal'

in other words, the variables  $Y_1, \dots, Y_n$  are required to be **uncorrelated**

since  $\mathbf{C}_X$  is symmetric, it has the decomposition:

$$\mathbf{C}_X = \mathbf{U} \mathbf{D} \mathbf{U}^T$$

where

- $\mathbf{D}$  is diagonal and its entries are eigenvalues of  $\mathbf{C}_X$
- $\mathbf{U}$  is unitary and the columns of  $\mathbf{U}$  are eigenvectors of  $\mathbf{C}_X$

**diagonalization:** pick  $\mathbf{A} = \mathbf{U}^T$  to obtain

$$\mathbf{C}_Y = \mathbf{A} \mathbf{C}_X \mathbf{A}^T = \mathbf{A} \mathbf{U} \mathbf{D} \mathbf{U}^T \mathbf{A}^T = \mathbf{U}^T \mathbf{U} \mathbf{D} \mathbf{U}^T \mathbf{U} = \mathbf{D}$$

as desired

one can write  $\mathbf{X}$  in terms of  $\mathbf{Y}$  as

$$\mathbf{X} = \mathbf{U}\mathbf{U}^T\mathbf{X} = \mathbf{U}\mathbf{Y} = \begin{bmatrix} U_1 & U_2 & \cdots & U_n \end{bmatrix} \begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix} = \sum_{k=1}^n Y_k U_k$$

this equation is called **Karhunen-Loève expansion**

- $\mathbf{X}$  can be expressed as a weighted sum of the eigenvectors  $U_k$
- the weighting coefficients are *uncorrelated* random variables  $Y_k$

**example:**  $\mathbf{X}$  has the covariance matrix  $\begin{bmatrix} 4 & 2 \\ 2 & 4 \end{bmatrix}$

design a transformation  $\mathbf{Y} = \mathbf{A}\mathbf{X}$  s.t. the covariance of  $\mathbf{Y}$  is diagonal

the eigenvalues of  $\mathbf{C}_{\mathbf{X}}$  and the corresponding eigenvectors are

$$\lambda_1 = 6, \quad u_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \lambda_2 = 2, \quad u_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$u_1$  and  $u_2$  are orthogonal, so if we normalize  $u_k$  so that  $\|u_k\| = 1$  then

$$U = \begin{bmatrix} \frac{u_1}{\sqrt{2}} & \frac{u_2}{\sqrt{2}} \end{bmatrix} \text{ is unitary}$$

therefore,  $\mathbf{C}_{\mathbf{X}} = UDU^T$  where

$$U = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}, \quad D = \begin{bmatrix} 6 & 0 \\ 0 & 2 \end{bmatrix}$$

thus, if we choose  $\mathbf{A} = U^T$  then  $\mathbf{C}_{\mathbf{Y}} = D$  which is diagonal as desired

# Whitening transformation

we wish to find a transformation  $\mathbf{Y} = \mathbf{A}\mathbf{X}$  such that

$$\mathbf{C}_Y = I$$

- a white noise property: the covariance is the identity matrix
- all components in  $\mathbf{Y}$  are all **uncorrelated**
- the variances of  $Y_k$  are **normalized** to 1

from  $\mathbf{C}_Y = \mathbf{A}\mathbf{C}_X\mathbf{A}^T$  and use the eigenvalue decomposition in  $\mathbf{C}_X$

$$\mathbf{C}_Y = \mathbf{A}\mathbf{U}\mathbf{D}\mathbf{U}^T\mathbf{A}^T = \mathbf{A}\mathbf{U}\mathbf{D}^{1/2}\mathbf{D}^{1/2}\mathbf{U}^T\mathbf{A}^T$$

denote  $\mathbf{D}^{1/2}$  the square root of  $\mathbf{D}$  with  $\mathbf{D} \succeq 0$ , *i.e.*,  $\mathbf{D}^{1/2}\mathbf{D}^{1/2} = \mathbf{D}$

$$\mathbf{D} = \mathbf{diag}(d_1, \dots, d_n) \implies \mathbf{D}^{1/2} = \mathbf{diag}(\sqrt{d_1}, \dots, \sqrt{d_n})$$

can you find  $\mathbf{A}$  that makes  $\mathbf{C}_Y$  the identity matrix ?

## Gaussian random vector

$X_1, \dots, X_n$  are said to be **jointly Gaussian** if their joint pdf is given by

$$f(\mathbf{x}) \triangleq f_{\mathbf{X}}(x_1, x_2, \dots, x_n) = \frac{1}{(2\pi)^{n/2} \det(\Sigma)^{1/2}} \exp -\frac{1}{2}(\mathbf{x}-\mu)^T \Sigma^{-1}(\mathbf{x}-\mu)$$

$\mu$  is the mean ( $n \times 1$ ) and  $\Sigma \succ 0$  is the covariance matrix ( $n \times n$ ):

$$\mu = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_n \end{bmatrix}, \quad \Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} & \cdots & \Sigma_{1n} \\ \Sigma_{21} & \Sigma_{22} & \cdots & \Sigma_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \Sigma_{n1} & \Sigma_{n2} & \cdots & \Sigma_{nn} \end{bmatrix}$$

and

$$\mu_k = \mathbf{E}[X_k], \quad \Sigma_{ij} = \mathbf{E}[(X_i - \mu_i)(X_j - \mu_j)]$$

**example:** the joint density function of  $\mathbf{x}$  (not normalized) is given by

$$f(x_1, x_2, x_3) = \exp - \frac{x_1^2 + 3x_2^2 + 2(x_3 - 1)^2 + 2x_1(x_3 - 1)}{2}$$

- $f$  is an exponential of *negative quadratic* in  $\mathbf{x}$  so  $\mathbf{x}$  must be a Gaussian

$$f(x_1, x_2, x_3) = \exp - \frac{1}{2} \begin{bmatrix} x_1 \\ x_2 \\ x_3 - 1 \end{bmatrix}^T \begin{bmatrix} 1 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 - 1 \end{bmatrix}$$

- the mean vector is  $(0, 0, 1)$  and the covariance matrix is


$$\mathbf{C} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 2 & 0 & -1 \\ 0 & 1/3 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

- the variance of  $x_1$  is highest while  $x_2$  is smallest
- $x_1$  and  $x_2$  are uncorrelated, so are  $x_2$  and  $x_3$

examples of Gaussian density contour (the exponent of exponential)


$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12} & \Sigma_{22} \end{bmatrix}^{-1} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 1$$

uncorrelated




$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

different variance



$$\Sigma = \begin{bmatrix} 1/2 & 0 \\ 0 & 1 \end{bmatrix}$$

correlated



$$\Sigma = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$$



# Properties of Gaussian variables

many results on Gaussian RVs can be obtained analytically:

- marginal's of  $\mathbf{X}$  is also Gaussian
- conditional pdf of  $X_k$  given the other variables is a Gaussian distribution
- uncorrelated Gaussian random variables are *independent*
- any affine transformation of a Gaussian is also a Gaussian

these are well-known facts

and more can be found in the areas of estimation, statistical learning, etc.

## Characteristic function of Gaussian

$$\Phi(\omega) = \Phi(\omega_1, \omega_2, \dots, \omega_n) = e^{j\mu^T \omega} e^{-\frac{\omega^T \Sigma \omega}{2}}$$

*Proof.* By definition and arranging the quadratic term in the power of exp

$$\begin{aligned}\Phi(\omega) &= \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} \int_{\mathbf{x}} e^{j\mathbf{x}^T \omega} e^{-\frac{(\mathbf{x}-\mu)^T \Sigma^{-1} (\mathbf{x}-\mu)}{2}} \mathbf{d}\mathbf{x} \\ &= \frac{e^{j\mu^T \omega} e^{-\frac{\omega^T \Sigma \omega}{2}}}{(2\pi)^{n/2} |\Sigma|^{1/2}} \int_{\mathbf{x}} e^{-\frac{(\mathbf{x}-\mu-j\Sigma\omega)^T \Sigma^{-1} (\mathbf{x}-\mu-j\Sigma\omega)}{2}} \mathbf{d}\mathbf{x} \\ &= \exp(j\mu^T \omega) \exp\left(-\frac{1}{2} \omega^T \Sigma \omega\right)\end{aligned}$$

(the integral equals 1 since it is a form of Gaussian distribution)

for one-dimensional Gaussian with zero mean and variance  $\Sigma = \sigma^2$ ,

$$\Phi(\omega) = e^{-\frac{\sigma^2 \omega^2}{2}}$$

## Affine Transformation of a Gaussian is Gaussian

let  $\mathbf{X}$  be an  $n$ -dimensional Gaussian,  $X \sim \mathcal{N}(\mu, \Sigma)$  and define

$$\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$$

where  $\mathbf{A}$  is  $m \times n$  and  $\mathbf{b}$  is  $m \times 1$  (so  $\mathbf{Y}$  is  $m \times 1$ )

$$\begin{aligned}\Phi_{\mathbf{Y}}(\omega) &= \mathbf{E}[e^{j\omega^T \mathbf{Y}}] = \mathbf{E}[e^{j\omega^T (\mathbf{A}\mathbf{X} + \mathbf{b})}] \\ &= \mathbf{E}[e^{j\omega^T \mathbf{A}\mathbf{X}} \cdot e^{j\omega^T \mathbf{b}}] = e^{j\omega^T \mathbf{b}} \Phi_{\mathbf{X}}(\mathbf{A}^T \omega) \\ &= e^{j\omega^T \mathbf{b}} \cdot e^{j\mu^T \mathbf{A}^T \omega} \cdot e^{-\omega^T \mathbf{A} \Sigma \mathbf{A}^T \omega / 2} \\ &= e^{j\omega^T (\mathbf{A}\mu + \mathbf{b})} \cdot e^{-\omega^T \mathbf{A} \Sigma \mathbf{A}^T \omega / 2}\end{aligned}$$

we read off that  $\mathbf{Y}$  is Gaussian with mean  $\mathbf{A}\mu + \mathbf{b}$  and covariance  $\mathbf{A}\Sigma\mathbf{A}^T$

## Marginal of Gaussian is Gaussian

the  $k^{\text{th}}$  component of  $\mathbf{X}$  is obtained by

$$X_k = [0 \quad \cdots \quad 1 \quad 0] \mathbf{X} \triangleq \mathbf{e}_k^T \mathbf{X}$$

( $\mathbf{e}_k$  is a standard unit column vector; all entries are zero except the  $k^{\text{th}}$  position)

hence,  $X_k$  is simply a linear transformation (in fact, a projection) of  $\mathbf{X}$

$X_k$  is then a Gaussian with mean

$$\mathbf{e}_k^T \boldsymbol{\mu} = \mu_k$$

and covariance

$$\mathbf{e}_k^T \boldsymbol{\Sigma} \mathbf{e}_k = \Sigma_{kk}$$

## Uncorrelated Gaussians are independent

suppose  $(\mathbf{X}, \mathbf{Y})$  is a jointly Gaussian vector with

$$\text{mean } \mu = \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} \quad \text{and covariance } \begin{bmatrix} \mathbf{C}_X & 0 \\ 0 & \mathbf{C}_Y \end{bmatrix}$$

in otherwords,  $X$  and  $Y$  are *uncorrelated* Gaussians:

$$\text{cov}(X, Y) = \mathbf{E}[XY^T] - \mathbf{E}[X] \mathbf{E}[Y]^T = 0$$

the joint density can be written as

$$\begin{aligned} f_{\mathbf{XY}}(\mathbf{x}, \mathbf{y}) &= \frac{1}{(2\pi)^n |\mathbf{C}_X|^{1/2} |\mathbf{C}_Y|^{1/2}} \exp -\frac{1}{2} \begin{bmatrix} \mathbf{x} - \mu_x \\ \mathbf{y} - \mu_y \end{bmatrix}^T \begin{bmatrix} \mathbf{C}_X^{-1} & 0 \\ 0 & \mathbf{C}_Y^{-1} \end{bmatrix} \begin{bmatrix} \mathbf{x} - \mu_x \\ \mathbf{y} - \mu_y \end{bmatrix} \\ &= \frac{1}{(2\pi)^{n/2} |\mathbf{C}_X|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\mu_x)^T \mathbf{C}_X^{-1}(\mathbf{x}-\mu_x)} \cdot \frac{1}{(2\pi)^{n/2} |\mathbf{C}_Y|^{1/2}} e^{-\frac{1}{2}(\mathbf{y}-\mu_y)^T \mathbf{C}_Y^{-1}(\mathbf{y}-\mu_y)} \end{aligned}$$

proving the independence

we can also see from the characteristic function

$$\begin{aligned}\Phi(\omega_1, \omega_2) &= \mathbf{E} \left[ \exp \left( j \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix}^T \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \right) \right] \\ &= \exp \left( j \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix}^T \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} \right) \cdot \exp - \frac{1}{2} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix}^T \begin{bmatrix} \mathbf{C}_X & 0 \\ 0 & \mathbf{C}_Y \end{bmatrix} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} \\ &= \exp (j\omega_1^T \mu_x) \exp - \frac{\omega_1^T \mathbf{C}_X \omega_1}{2} \cdot \exp (j\omega_2^T \mu_y) \cdot \exp - \frac{\omega_2^T \mathbf{C}_Y \omega_2}{2} \\ &\triangleq \Phi_1(\omega_1) \cdot \Phi_2(\omega_2)\end{aligned}$$

proving the independence

## Conditional of Gaussian is Gaussian

let  $\mathbf{Z}$  be an  $n$ -dimensional Gaussian which can be decomposed as

$$\mathbf{Z} = \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{xy}^T & \Sigma_{yy} \end{bmatrix} \right)$$

the conditional pdf of  $\mathbf{X}$  given  $\mathbf{Y}$  is also Gaussian with conditional mean

$$\mu_{\mathbf{X}|\mathbf{Y}} = \mu_x + \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{Y} - \mu_y)$$

and conditional covariance

$$\Sigma_{\mathbf{X}|\mathbf{Y}} = \Sigma_x - \Sigma_{xy} \Sigma_{yy}^{-1} \Sigma_{xy}^T$$

**Proof:**

from the **matrix inversion lemma**,  $\Sigma^{-1}$  can be written as

$$\Sigma^{-1} = \begin{bmatrix} S^{-1} & -S^{-1}\Sigma_{xy}\Sigma_{yy}^{-1} \\ -\Sigma_{yy}^{-1}\Sigma_{xy}^T S^{-1} & \Sigma_{yy}^{-1} + \Sigma_{yy}^{-1}\Sigma_{xy}^T S^{-1}\Sigma_{xy}\Sigma_{yy}^{-1} \end{bmatrix}$$

where  $S$  is called the **Schur complement of  $\Sigma_{xx}$  in  $\Sigma$**  and

$$\begin{aligned} S &= \Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{xy}^T \\ \det \Sigma &= \det S \cdot \det \Sigma_{yy} \end{aligned}$$

we can show that  $\Sigma \succ 0$  if and only if  $S \succ 0$  and  $\Sigma_{yy} \succ 0$



from  $f_{\mathbf{X}|\mathbf{Y}}(\mathbf{x}|\mathbf{y}) = f_{\mathbf{X}}(\mathbf{x}, \mathbf{y})/f_{\mathbf{Y}}(\mathbf{y})$ , we calculate the exponent terms

$$\begin{aligned}
 & \begin{bmatrix} \mathbf{x} - \mu_x \\ \mathbf{y} - \mu_y \end{bmatrix}^T \Sigma^{-1} \begin{bmatrix} \mathbf{x} - \mu_x \\ \mathbf{y} - \mu_y \end{bmatrix} - (\mathbf{y} - \mu_y)^T \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y) \\
 &= (\mathbf{x} - \mu_x)^T S^{-1} (\mathbf{x} - \mu_x) - (\mathbf{x} - \mu_x)^T S^{-1} \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y) \\
 &\quad - (\mathbf{y} - \mu_y)^T \Sigma_{yy}^{-1} \Sigma_{xy}^T S^{-1} (\mathbf{x} - \mu_x) \\
 &\quad + (\mathbf{y} - \mu_y)^T (\Sigma_{yy}^{-1} \Sigma_{xy}^T S^{-1} \Sigma_{xy} \Sigma_{yy}^{-1}) (\mathbf{y} - \mu_y) \\
 &= [\mathbf{x} - \mu_x - \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y)]^T S^{-1} [\mathbf{x} - \mu_x - \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y)] \\
 &\triangleq (\mathbf{x} - \mu_{\mathbf{X}|\mathbf{Y}})^T \Sigma_{\mathbf{X}|\mathbf{Y}}^{-1} (\mathbf{x} - \mu_{\mathbf{X}|\mathbf{Y}})
 \end{aligned}$$

$f_{\mathbf{X}|\mathbf{Y}}(\mathbf{x}|\mathbf{y})$  is an exponential of quadratic function in  $\mathbf{x}$

so it has a form of Gaussian

## Standard Gaussian vectors

for an  $n$ -dimensional Gaussian vector  $X \sim \mathcal{N}(\mu, \mathbf{C})$  with  $\mathbf{C} \succ 0$

let  $\mathbf{A}$  be an  $n \times n$  invertible matrix such that

$$\mathbf{A}\mathbf{A}^T = \mathbf{C}$$

( $\mathbf{A}$  is called a **factor** of  $\mathbf{C}$ )

then the random vector

$$\mathbf{Z} = \mathbf{A}^{-1}(\mathbf{X} - \mu)$$

is a standard Gaussian vector, *i.e.*,

$$\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

(obtain  $\mathbf{A}$  via eigenvalue decomposition or Cholesky factorization)

# Functions of random vectors

- minimum and maximum of random variables
- general transformation
- affine transformation

## Minimum and Maximum of RVs

let  $X_1, X_2, \dots, X_n$  be independent RVs

define the minimum and maximum of RVs by

$$Y = \min(X_1, X_2, \dots, X_n), \quad Z = \max(X_1, X_2, \dots, X_n)$$

the maximum of  $X_1, X_2, \dots, X_n$  is less than  $z$  iff  $X_i \leq z$  for all  $i$ , so

$$F_Z(z) = P(X_1 \leq z)P(X_2 \leq z) \cdots P(X_n \leq z) = (F_X(z))^n$$


the minimum of  $X_1, X_2, \dots, X_n$  is greater than  $y$  iff  $X_i \geq y$  for all  $i$ , so

$$1 - F_Y(y) = P(X_1 > y)P(X_2 > y) \cdots P(X_n > y) = (1 - F_X(y))^n$$

and

$$F_Y(y) = 1 - (1 - F_X(y))^n$$

# Example: Merging of independent Poisson arrivals



- $T_i$  denotes the interarrival times for source  $i$
- $T_i$  has exponential distribution with rate  $\lambda_i$
- find the distribution of the interarrival times between consecutive requests at server

each  $T_i$  satisfies the memoryless property, so the time that has elapsed since the last arrival is irrelevant

the time until the next arrival at the multiplexer is

$$Z = \min(T_1, T_2, \dots, T_n)$$

therefore, the cdf of  $Z$  can be computed by:

$$\begin{aligned} 1 - F_Z(z) &= P(\min(T_1, T_2, \dots, T_n) > z) \\ &= P(T_1 > z)P(T_2 > z) \cdots P(T_n > z) \\ &= (1 - F_{T_1}(z))(1 - F_{T_2}(z)) \cdots (1 - F_{T_n}(z)) \\ &= e^{-\lambda_1 z} \cdots e^{-\lambda_n z} = e^{-(\lambda_1 + \lambda_2 + \cdots + \lambda_n)z} \end{aligned}$$

the interarrival time is an exponential RV with rate  $\lambda_1 + \lambda_2 + \cdots + \lambda_n$

## General transformation

let  $\mathbf{X}$  be a vector random variable

define  $\mathbf{Z} = g(\mathbf{X}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$  and assume that  $g$  is invertible

so that for  $\mathbf{Z} = \mathbf{z}$  we can solve for  $\mathbf{x}$  uniquely:

$$\mathbf{x} = g^{-1}(\mathbf{z})$$

then the joint pdf of  $\mathbf{Z}$  is given by

$$f_{\mathbf{Z}}(\mathbf{z}) = \frac{f_{\mathbf{X}}(g^{-1}(\mathbf{z}))}{|\det J|}$$

where  $\det J$  is the determinant of the Jacobian matrix:

$$J = \begin{bmatrix} \partial g_1 / \partial x_1 & \cdots & \partial g_1 / \partial x_n \\ \vdots & \ddots & \vdots \\ \partial g_n / \partial x_1 & \cdots & \partial g_n / \partial x_n \end{bmatrix}$$

## Affine transformation

if  $\mathbf{X}$  is a continuous random vector and  $\mathbf{A}$  is an invertible matrix  
then  $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$  has pdf

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{1}{|\det(\mathbf{A})|} f_{\mathbf{X}}(\mathbf{A}^{-1}(\mathbf{y} - \mathbf{b}))$$

**Gaussian case:** let  $\mathbf{X} \sim \mathcal{N}(0, \Sigma)$

$$\begin{aligned} f_{\mathbf{Y}}(\mathbf{y}) &= \frac{1}{(2\pi)^{n/2} |\det \mathbf{A}| |\Sigma|^{1/2}} \exp -\frac{1}{2}(\mathbf{y} - \mathbf{b})^T \mathbf{A}^{-T} \Sigma^{-1} \mathbf{A}^{-1}(\mathbf{y} - \mathbf{b}) \\ &= \frac{1}{(2\pi)^{n/2} |\mathbf{A}\Sigma\mathbf{A}^T|^{1/2}} \exp -\frac{1}{2}(\mathbf{y} - \mathbf{b})^T (\mathbf{A}\Sigma\mathbf{A}^T)^{-1}(\mathbf{y} - \mathbf{b}) \end{aligned}$$

we read off that  $\mathbf{Y}$  is also Gaussian with mean  $\mathbf{b}$  and covariance  $\mathbf{A}\Sigma\mathbf{A}^T$

this agrees with the result in page 5-16 and 5-27



## Example: Sum of jointly Gaussian

a special case of linear transformation is

$$Z = a_1X_1 + a_2X_2 + \cdots + a_nX_n$$

where  $X_1, \dots, X_n$  are jointly Gaussian

$Z$  can be written as

$$Z = [a_1 \quad \cdots \quad a_n] \begin{bmatrix} X_1 \\ \vdots \\ X_n \end{bmatrix} \triangleq \mathbf{A}\mathbf{X}$$

$Z$  is simply a linear transformation of a Gaussian

therefore,  $Z$  is Gaussian with mean

$$\mathbf{E}[Z] = \mathbf{A}\boldsymbol{\mu} = \sum_{i=1} a_i \mathbf{E}[X_i]$$

and variance

$$\mathbf{var}(Z) = \mathbf{cov}(Z) = \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T = \sum_{i=1}^n \sum_{j=1}^n a_i a_j \mathbf{cov}(X_i, X_j)$$

if  $X_1, \dots, X_n$  are *independent* Gaussian, i.e.,

$$\mathbf{cov}(X_i, X_j) = 0$$

then the variance of  $Z$  is reduced to

$$\mathbf{var}(Z) = \sum_{i=1}^n a_i^2 \mathbf{cov}(X_i, X_i) = \sum_{i=1}^n a_i^2 \mathbf{var}(X_i)$$

# References

Chapter 6 in

A. Leon-Garcia, *Probability, Statistics, and Random Processes for Electrical Engineering*, 3rd edition, Pearson Prentice Hall, 2009